

Application No. 10/675,470

**Amendments to the Claims:**

This listing of claims will replace all prior versions, and listings, of claims in the application:

**Listing of Claims:**

1-7. (Canceled)

8. (Currently Amended) A method of inhibiting angiogenesis in pathological conditions where increased angiogenesis and coincidental vascular perfusion are clinically detrimental, comprising the steps of: producing an AT<sub>4</sub> receptor ligand, having a structure selected from the group consisting of ~~with the structure~~  $\text{NH}_3^+$ -norleucine-tyrosine-isoleucine-histidine-COO<sup>-</sup>, and norleucine-tyrosine-isoleucine-(6-amino-hexanoic acid)-CONH<sub>2</sub>; or ~~norleucine-tyrosine-leucine- $\text{I}^{\text{L}}$ -(CH<sub>2</sub>-HN<sub>2</sub>)<sup>3-4</sup>-histidine-proline-phenylalanine)-COO<sup>-</sup>~~, and administering the AT<sub>4</sub> receptor ligand.

9. (Currently Amended) The method of inhibiting angiogenesis ~~accordingly~~ according to claim 8 or claim 29, further comprising the delivery of the AT<sub>4</sub> receptor ligand locally.

10. (Currently Amended) The method of inhibiting angiogenesis according to claim 8 or claim 29, further comprising the delivery of the AT<sub>4</sub> receptor ligand intravascularly.

11. (Currently Amended) The method of inhibiting angiogenesis according to claim 8 or claim 29, further comprising the delivery of the AT<sub>4</sub> receptor ligand intramuscularly.

12. (Currently Amended) The method of inhibiting angiogenesis according to claim 8 or claim 29, further comprising the delivery of the AT<sub>4</sub> receptor ligand intraperitoneally.

Application No. 10/675,470

Best Available Copy

13. (Currently Amended) The method of inhibiting angiogenesis according to claim 8 or claim 29, further comprising the delivery of the AT<sub>4</sub> receptor ligand subcutaneously.
14. (Currently Amended) The method of inhibiting angiogenesis according to claim 8 or claim 29, further comprising the delivery of the AT<sub>4</sub> receptor ligand orally.
15. (Currently Amended) A method of inhibiting the growth and metastasis of solid tumors, comprising the steps of: producing an AT<sub>4</sub> receptor ligand, having a structure selected from the group consisting of: ~~with the structure~~ NH<sub>3</sub><sup>+</sup>-norleucine-tyrosine-isoleucine-histidine-COO<sup>-</sup>, and norleucine-tyrosine-isoleucine-(6-amino-hexanoic acid)-CONH<sub>2</sub>; ~~or norleucine-tyrosine-leucine-N-(CH<sub>2</sub>-HN<sub>2</sub>)<sup>3-4</sup>-histidine-proline-phenylalanine)-COO<sup>-</sup>~~ and administering the AT<sub>4</sub> receptor ligand.
16. (Currently Amended) The method of inhibiting the growth and metastasis of solid tumors according to claim 15 or claim 30, further comprising delivery of the AT<sub>4</sub> receptor ligand locally.
17. (Currently Amended) The method of inhibiting the growth and metastasis of solid tumors according to claim 15 or claim 30, further comprising the delivery of the AT<sub>4</sub> receptor ligand intravascularly.
18. (Currently Amended) The method of inhibiting the growth and metastasis of solid tumors according to claim 15 or claim 30, further comprising the delivery of the AT<sub>4</sub> receptor ligand intramuscularly.
19. (Currently Amended) The method of inhibiting the growth and metastasis of solid tumors according to claim 15 or claim 30, further comprising the delivery of the AT<sub>4</sub> receptor ligand intraperitoneally.

Application No. 10/675,470

20. (Currently Amended) The method of inhibiting the growth and metastasis of solid tumors according to claim 15 or claim 30, further comprising the step of applying the AT<sub>4</sub> receptor ligand subcutaneously.

21. (Currently Amended) The method of inhibiting the growth and metastasis of solid tumors according to claim 15 or claim 30, further comprising the step of applying the AT<sub>4</sub> receptor ligand orally.

22. (Currently Amended) A method of inhibiting the growth and metastasis of breast cancer, comprising the steps of: producing an AT<sub>4</sub> receptor ligand, having a structure selected from the group consisting of: with the structure NH<sub>3</sub><sup>+</sup>-norleucine-tyrosine-isoleucine-histidine-COO<sup>-</sup>, and norleucine-tyrosine-isoleucine-(6-amino-hexanoic acid)-CONH<sub>2</sub>; or ~~norleucine-tyrosine-leucine-W-(CH<sub>2</sub>-HN<sub>2</sub>)<sup>3-4</sup>-histidine-proline-phenylalanine)-COO<sup>-</sup>~~ and administering the AT<sub>4</sub> receptor ligand.

23. (Currently Amended) The method of inhibiting the growth and metastasis of breast cancer according to claim 22 or claim 31, further comprising the delivery of the AT<sub>4</sub> receptor ligand locally to the tumor.

24. (Currently Amended) The method of inhibiting the growth and metastasis of breast cancer according to claim 22 or claim 31, further comprising the delivery of the AT<sub>4</sub> receptor ligand intravascularly.

25. (Currently Amended) The method of inhibiting the growth and metastasis of breast cancer according to claim 22 or claim 31, further comprising the delivery of the AT<sub>4</sub> receptor ligand intramuscularly.

26. (Currently Amended) The method of inhibiting the growth and metastasis of breast cancer according to claim 22 or claim 31, further comprising the delivery of the AT<sub>4</sub> receptor ligand intraperitoneally.

Application No. 10/675,470

27. (Currently Amended) The method of inhibiting the growth and metastasis of breast cancer according to claim 22 or claim 31, further comprising the delivery of the AT<sub>4</sub> receptor ligand subcutaneously.

28. (Currently Amended) The method of inhibiting the growth and metastasis of breast cancer according to claim 22 or claim 31, further comprising the delivery of the AT<sub>4</sub> receptor ligand orally.

29. (New) A method of inhibiting angiogenesis in pathological conditions where increased angiogenesis and coincidental vascular perfusion are clinically detrimental, comprising the steps of: producing an AT<sub>4</sub> receptor ligand having a structure of norleucine-tyrosine-leucine-Ψ-(CH<sub>2</sub>-NH<sub>2</sub>)<sup>3-4</sup>-histidine-proline-phenylalanine-COO<sup>-</sup>; and administering the AT<sub>4</sub> receptor ligand.

30. (New) A method of inhibiting the growth and metastasis of solid tumors, comprising the steps of: producing an AT<sub>4</sub> receptor ligand having a structure of: norleucine-tyrosine-leucine-Ψ-(CH<sub>2</sub>-NH<sub>2</sub>)<sup>3-4</sup>-histidine-proline-phenylalanine-COO<sup>-</sup>; and administering the AT<sub>4</sub> receptor ligand.

31. (New) A method of inhibiting the growth and metastasis of breast cancer, comprising the steps of: producing an AT<sub>4</sub> receptor ligand having a structure of: norleucine-tyrosine-leucine-Ψ-(CH<sub>2</sub>-NH<sub>2</sub>)<sup>3-4</sup>-histidine-proline-phenylalanine-COO<sup>-</sup>; and administering the AT<sub>4</sub> receptor ligand.

Best Available Copy